



kathy.mckay@bullpg.com

sven.magnussen@bullpg.com



COMPACT

- Most compact series
- Lots of blooms and large flowers
- 12-14" height
- Great for 6" gallon, 8" whiskey barrel



Mia



Emma



Marilyn

MEDIUM

- Medium sized habit
- Large flowers and blooms
- 14-16" height
- Great for 6" gallon, 8-10" whiskey barrel





Carla

Carolina









Lucia

Maya

Ophelia

Sarah

GRANDE

- Largest size and blooms of the series 15-18" height
- Some over 6" in diameter

- 8", 10", 12" pots







Juliana

Laura

Paulina







GENERAL INFORMATION

Rudbeckia Sunbeckia series is a hirta hybrid that was bred in Germany and is known for its large flower size and long-lasting blooms. It is available as tissue culture and is recommended to be produced in a 72 or 105 cell tray (app 5-6 week propagation time). It is also available as a liner.

Due to its strong vigor it has a short crop time of 8-10 weeks after transplanting. The Sunbeckia series requires long day treatment. Without this period it is not advised to start transplanting prior to week 12. In addition to being a spring crop, we recommend continue transplanting in a 3 weeks rhythm. The start in week 18 till 26 results in perfect summer and fall production. A flowering 'Sunbeckia' has outstanding garden performance as well as great in a patio container.

TRANSPLANTING

Remove liner carefully from trays so as to not damage roots. Plant liner even with soil line when transplanting. Keep soil surface of liner level with surface of finished pot.

CONTAINER SIZES AND CROP TIMES

- Gallon/6" 1ppp. 8-10 weeks from transplant.
- 8": 1ppp. 10-11 weeks from transplant.

GROWING MEDIA

Plant into a well-drained soil mix with a pH range of 5.5-6.0. Soil moisture should contain coarse components for good drainage.

TEMPERATURE

Day temperature: 60-64°F (15-18°C) Night Temperature: 50-55°F (10-13°C) Vernalization is not require for flowering

LIGHT

During propagation, it is important to have short days (10 hours or less). After transplanting, long days are required for flowering (minimum 12 hours; recommended 14 hours).

WATER AND FEED

Rudbeckia Sunbeckia should be grown using a well-balanced fertilizer (15-15-15 or similar) approximately once a week. Calcium nitrate can help with growth if plants are too small. 200-250 ppm N is recommended.

It is important to maintain consistent moisture in the soil. Overhead irrigation or drip irrigation is best.

GROWTH REGULATORS

PGR can be applied when the bottom shoots start to stretch. Also, later when the flower buds are visible and stems start to stretch. B-Nine is beneficial and can be applied approximately 3-4 weeks after transplant.

If needed, Bonzi in the later weeks can be applied.

INSECTS AND DISEASES

Mites, Thrips and Aphids.
Botrytis, Pythium and Phytophthora.

